

UNCLASSIFIED

AD NUMBER	
AD370835	
CLASSIFICATION CHANGES	
TO:	unclassified
FROM:	confidential
LIMITATION CHANGES	
TO:	Approved for public release, distribution unlimited
FROM:	Controlling DoD Organization: Office of Naval Research, Arlington, VA 22217.
AUTHORITY	
Office of Naval Research ltr dtd 4 May 1977; Office of Naval Research ltr dtd 4 May 1977	

THIS PAGE IS UNCLASSIFIED

# **SECURITY**

---

# **MARKING**

**The classified or limited status of this report applies to each page, unless otherwise marked.**

**Separate page printouts MUST be marked accordingly.**


---

**THIS DOCUMENT CONTAINS INFORMATION AFFECTING THE NATIONAL DEFENSE OF THE UNITED STATES WITHIN THE MEANING OF THE ESPIONAGE LAWS, TITLE 18, U.S.C., SECTIONS 793 AND 794. THE TRANSMISSION OR THE REVELATION OF ITS CONTENTS IN ANY MANNER TO AN UNAUTHORIZED PERSON IS PROHIBITED BY LAW.**

**NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.**

**CONFIDENTIAL**

370835




**ROCKETDYNE**

A DIVISION OF NORTH AMERICAN AVIATION, INC.

THIS MATERIAL CONTAINS INFORMATION AFFECTING THE NATIONAL DEFENSE OF THE UNITED STATES WITHIN THE MEANING OF THE ESPIONAGE LAWS, TITLE 18 U.S.C., SECTIONS 793 AND 794, THE TRANSMISSION OR REVELATION OF WHICH IN ANY MANNER TO AN UNAUTHORIZED PERSON IS PROHIBITED BY LAW.

**CONFIDENTIAL**



**CONFIDENTIAL**

**ROCKETDYNE**  
A DIVISION OF NORTH AMERICAN AVIATION INC.  
3633 CANOGA AVENUE, CANOGA PARK, CALIFORNIA

R-5883-6

(Unclassified Title)

QUARTERLY PROGRESS REPORT,  
INORGANIC HALOGEN OXIDIZERS  
(29 November 1965 through 28 February 1966)

Group 4  
Downgraded at 3-Year Intervals  
Declassified After 12 Years

Contract Nonr 4428(00)

Sponsored by Advanced Research Projects Agency  
Washington 25, D.C.  
ARPA Order No. 23

**PREPARED BY**

D. Pilipovich  
H. F. Bauer  
C. J. Schack  
C. B. Lindahl

Reproduction in whole or in  
part is permitted for any  
purpose of the  
United States Government

**APPROVED BY**

*J. Silverman*  
J. Silverman  
Section Chief  
Chemistry

Research Division  
**REVISIONS**

NO. OF PAGES 22 & v

DATE 30 March 1966

DATE	REV. BY	PAGES AFFECTED	REMARKS

FORM R18-G-1 REV 4-64

**CONFIDENTIAL**

This material contains information affecting the national defense of the United States within the meaning of the Espionage Laws, Title 18, U.S.C., Sections 793 and 794, the transmission or revelation of which in any manner to an unauthorized person is prohibited by law.

NOTICE

UNITED STATES PATENT OFFICE SECRECY ORDER

A patent application has been filed in the U. S. Patent Office by North American Aviation, Inc. based upon subject matter included herein or related hereto, and the Secrecy Order appended hereto has been issued thereon pursuant to Title 35, United States Code (1952) Sections 181-188. Further dissemination of said subject matter is prohibited except in strict compliance with said order. The recipient of this document is requested to notify all persons who will have access to this material of the Secrecy Order. Penalties for violation of a Secrecy Order include a fine of up to \$10,000 or imprisonment for not more than two years, or both.

DEPARTMENT OF COMMERCE  
United States Patent Office  
Washington

DEPARTMENT OF COMMERCE  
United States Patent Office  
Washington

SECRECY ORDER

NOTICE: To the applicant above named, his heirs, and any and all his assignees, attorneys and agents, hereinafter designated principals:

You are hereby notified that your application as above identified has been found to contain subject matter, the unauthorized disclosure of which might be detrimental to the public safety or defense, and you are ordered in nowise to publish or disclose the invention or any material information with respect thereto, including hitherto unpublished details of the subject matter of said application, in any way to any person not cognizant of the invention prior to the date of the order, including any employee of the principals, but to keep the same secret except by written permission first obtained of the Commissioner of Patents, under the penalties of 35 U. S. C. (1952) 182, 186.

Any other application which contains any significant part of the subject matter of the above identified application falls within the scope of this order. If such other application does not stand under a secrecy order, it and the common subject matter should be brought to the attention of the Patent Security Division, Patent Office.

If prior to the issuance of the secrecy order any significant part of the subject matter has been revealed to any person, the principals shall promptly inform such person of the secrecy order and the penalties for improper disclosure.

This order should not be construed in any way to mean that the Government has adopted or contemplates adoption of the alleged invention disclosed in this application, nor is it any indication of the value of such invention.

PERMIT FOR PATENT APPLICATION  
CLASSIFIED BY GOVERNMENT CONTRACT

This permit authorizes the principals, as designated in the secrecy order, to take any action with regard to the subject matter of the application, to the extent authorized by the security requirements of the Government contract which imposes the highest security classification on the subject matter of this application, except that this permit does not authorize export of this application or the subject matter thereof, whether for the foreign filing of corresponding applications or otherwise. Permission for such export must be specifically obtained from the Patent Office.

**CONFIDENTIAL**



**ROCKETDYNE • A DIVISION OF NORTH AMERICAN AVIATION, INC.**

**FOREWORD**

The research reported herein was supported by the Advanced Research Projects Agency through the Office of Naval Research, Power Branch, Code 429, with Mr. Richard L. Hanson as Scientific Officer. This report was prepared in compliance with Section H of Contract Nonr 4428(00) under ARPA Order No. 23, and covers the period 29 November 1965 through 28 February 1966. This work was carried out in the Synthetic Chemistry Group with Dr. E. A. Lawton as Group Scientist. Dr. D. Pilipovich provided the technical direction as Project Scientist. Full-time associates connected with the technical effort were Dr. H. F. Bauer, responsible for coordinating this report; Dr. C. J. Schack; and Dr. C. B. Lindahl.

**CONFIDENTIAL**

CONFIDENTIAL



ROCKETDYNE • A DIVISION OF NORTH AMERICAN AVIATION, INC.

ABSTRACT

Oxychlorine trifluoride, Florox, has been synthesized from  $\text{Cl}_2\text{O} + \text{F}_2$  in the presence of CsF, RbF, KF, and NaF as well as in their absence. Although the complexes of  $\text{Cl}_2\text{O}$  with CsF, RbF, and KF were probably involved, no substantial increase in conversion to Florox was obtained over the systems with uncomplexed  $\text{Cl}_2\text{O}$ . Oxychlorine trifluoride has also been synthesized from uncomplexed chlorine nitrate in 84-percent yield. Fluorination in situ of the  $\text{Cl}_2 + \text{HgO}$  reaction products has also produced  $\text{ClF}_3\text{O}$ .

Florox was stable in excess hydrogen fluoride or excess oxygen at ambient temperature and stable to 5-day storage at 71 C in Monel. In a flow system, thermal decomposition of Florox started near 300 C and was complete below 586 C. The predominant product recovered above 300 C was chlorine monofluoride.

The complex between Florox and CsF dissociated below 150 C and allowed complete removal of contaminant  $\text{ClF}_3$  as the remaining nonvolatile solid  $\text{CsClF}_4$ . The complex between  $\text{ClF}_3\text{O}$  and KF was less stable, dissociating slowly with pumping at ambient temperature. Investigation of the FNO complex of  $\text{ClF}_3\text{O}$  by  $\text{F}^{19}$  n.m.r. showed exchange at -77 C; an infrared study at -196 C indicated the complex to be predominantly covalent.

CONFIDENTIAL



ROCKETDYNE • A DIVISION OF NORTH AMERICAN AVIATION, INC.

## CONFIDENTIAL

Attempts to reduce  $\text{ClF}_3\text{O}$  to  $\text{FClO}$  have been unsuccessfully attempted with  $\text{AgF}$ ,  $\text{Cl}_2$ , and  $\text{Cl}_2\text{O}$ . Fluorination of  $\text{Cl}_2\text{O}$  by  $\text{F}_2$ ,  $\text{ClF}_3$ , and  $\text{ClF}_5$  has also failed to produce  $\text{FClO}$ . However, fluorination of the products of the chlorine-mercuric oxide reaction has produced some unknown, labeled Compound C, in small quantity. Some infrared evidence for traces of Compound C was also obtained in the thermal dissociation of  $\text{ClF}_3\text{O}$ .

As a new route to oxychlorine fluorides, the reported intermediate  $\text{ClO}\cdot\text{AsF}_5$  has been investigated. Characterization has been unsuccessful thus far, and the solid chlorine content from mass balance and analysis has been quite low.

The synthesis of new bromine oxyfluorides from  $\text{Br}_2\text{O}$  complexes failed, giving mainly  $\text{BrF}_5$ . Bromine nitrate has been prepared as a possible alternate precursor to  $\text{BrF}_3\text{O}$  or  $\text{BrF}_5\text{O}$ .

Although the nature of the bonding in the  $\text{Cl}_2\text{O}\cdot\text{CsF}$  complex has not been clarified, a reproducible composition,  $\text{CsF}\cdot 1.5 \text{ Cl}_2\text{O}$  has been obtained near  $-80^\circ \text{C}$ .

(Confidential Abstract)

## CONFIDENTIAL



# CONFIDENTIAL



ROCKETDYNE • A DIVISION OF NORTH AMERICAN AVIATION, INC.

## CONTENTS

Foreword . . . . .	ii
Abstract . . . . .	iii
Introduction . . . . .	1
Discussion . . . . .	2
Florox Studies . . . . .	2
Florox Complexes with Alkali Metal Fluorides . . . . .	11
Attempts at New Oxyhalogen Fluoride Oxidizers . . . . .	15
Cesium Fluoride-Chlorine Monoxide Complex Studies . . . . .	19
Experimental Effort . . . . .	20
Low-Temperature Infrared Cell . . . . .	20
Synthesis of Florox . . . . .	20
References . . . . .	22

CONFIDENTIAL

CONFIDENTIAL



ROCKETDYNE • A DIVISION OF NORTH AMERICAN AVIATION INC

#### INTRODUCTION

The work reported herein reflects effort during the past quarter in the general area of interhalogen fluorides and their derivatives. The laboratory studies encompassed roughly three areas of interest. One area involved the completion of previously initiated low-temperature fluorinations of  $\text{Cl}_2\text{O}$  and  $\text{ClONO}_2$  in Florox syntheses. These studies not only gave yield data of interest but also provided working quantities of  $\text{ClF}_3\text{O}$  for further chemical investigation.

The studies on  $\text{ClF}_3\text{O}$  are of a dual nature. One area is concerned with both the possible redox reaction of  $\text{ClF}_3\text{O}$  and the decomposition reactions which may lead to  $\text{FClO}$ . In addition, both infrared and n.m.r. examination of the complexes are of interest and are reported herein.

Finally, the preparation of Br, F, O species is being sought through the fluorination of  $\text{Br}_2\text{O}$  and  $\text{BrONO}_2$ . These studies are logical extensions in the area of interhalogen derivatives as dictated by the ease of formation of  $\text{ClF}_3\text{O}$ .

CONFIDENTIAL



## CONFIDENTIAL

## DISCUSSION

## FLOROX STUDIES

Florox Synthesis from  $\text{Cl}_2\text{O}$ 

The preparation of  $\text{ClF}_3\text{O}$  by fluorination of  $\text{Cl}_2\text{O}$  in the presence of alkali metal fluoride has been extended to NaF and RbF in addition to the previously reported results with KF and CsF (Ref. 1). In addition, preparation of  $\text{ClF}_3\text{O}$  in good yield has been achieved in the absence of added alkali metal fluoride. The results of preparative runs are presented in Table 1.

TABLE 1

SYNTHESIS OF FLOROX

Alkali Metal Fluoride	Percent Yield	Side Products
CsF	0 to 82	$\text{FClO}_2$ , $\text{ClF}_3$
RbF	> 25	$\text{FClO}_2$ , $\text{ClF}_3$
KF	43, 29	$\text{FClO}_2$ , $\text{ClF}_3$ , $\text{ClF}$
NaF	73, 81	$\text{FClO}_2$ , $\text{ClF}$
None	39, 63	$\text{FClO}_2$ , $\text{ClF}$ , $\text{ClF}_3$

The variation in yields caused by different added alkali metal fluorides is not necessarily significant because of the difference in yields in

CONFIDENTIAL

CONFIDENTIAL



ROCKETDYNE • A DIVISION OF NORTH AMERICAN AVIATION, INC.

apparently identical preparations (particularly during CsF experiments where the most data have been gained). Amounts of reactants and reaction times involved, and the partial decomposition of  $\text{Cl}_2\text{O}$  in metal reaction systems were kept substantially the same.

#### Synthesis of Florox from $\text{ClNO}_2$


The synthesis of Florox by fluorination of the CsF complex of  $\text{ClNO}_2$  was described in earlier reports (Ref. 1 and 2). It was of interest also to utilize  $\text{ClONO}_2$  as an intermediate for the preparation of  $\text{FClO}$ . Accordingly, a reaction was attempted using uncomplexed  $\text{ClNO}_2$  and  $\text{F}_2$  at  $-80^\circ\text{C}$ . After several days it was found that Florox was formed in 84-percent yield:



Thus, it has been demonstrated that the use of CsF was not essential and two steps of the previous reaction sequences, complexing and pyrolysis, to liberate  $\text{ClF}_3\text{O}$  were eliminated.

#### Synthesis from $\text{HgCl}_2\text{-Cl}_2\text{O}$

The synthesis of  $\text{ClF}_3\text{O}$  from  $\text{Cl}_2\text{O}$  in the past has required the vacuum transfer of  $\text{Cl}_2\text{O}$  from an apparent  $\text{Cl}_2\text{O}\cdot\text{HgCl}_2$  complex with warming to another reaction vessel. It has been found that if the solid products from the  $\text{HgO} + \text{Cl}_2$  reaction are fluorinated in situ,  $\text{ClF}_3\text{O}$  and  $\text{ClF}_3$  are produced. This is the first time Florox has been prepared in this manner:



CONFIDENTIAL

CONFIDENTIAL



ROCKETDYNE • A DIVISION OF NORTH AMERICAN AVIATION, INC.

Thermal Decomposition of Florox

Pyrolysis of the nonvolatile solid formed by low-temperature fluorination of the  $\text{Cl}_2\text{O}-\text{CsF}$  complex has, on at least three occasions, yielded traces of an unknown species thought to be  $\text{FClO}$  (Compound C). It was thought that Compound C might result from pyrolysis of  $\text{ClF}_3\text{O}$  as in Eq. 3.



Representative results of experiments where Florox at low pressure was passed through a Monel tube held at temperatures from 300 to 585 C are presented in Table 2. As is seen in Table 2, decomposition of  $\text{ClF}_3\text{O}$  does not occur to a large extent under these flow conditions at temperatures less than 400 C. Products found at 300 C were  $\text{ClF}_3$  and possibly

TABLE 2  
PYROLYSIS OF  $\text{ClF}_3\text{O}$

Temperature, C	Percent Florox Recovered	Condensable Product Formed at -196 C (as percent of Florox passed)	Products
300	--	~ 1.5	$\text{ClF}_3$ , $\text{FClO}_2$
400	99	2	$\text{ClF}$ , trace Compound C
450	82	16	$\text{ClF}$
500	46	54	$\text{ClF}$
500	38	62	$\text{FClO}_2$ , $\text{ClF}$
585	0	98	$\text{ClF}$ , trace Compound C

R-5883-6

CONFIDENTIAL

## CONFIDENTIAL



ROCKETDYNE • A DIVISION OF NORTH AMERICAN AVIATION INC

FC10<sub>2</sub>. At 400 C. small amounts of ClF as well as ClF<sub>3</sub> were found, and at higher temperatures ClF was the only product. Two reactions yielded traces of unknown material, possibly Compound C. Material balances agree with the expected evolution of one molecule of ClF (or ClF<sub>3</sub>) per molecule of decomposed ClF<sub>3</sub>O according to the overall reaction:



Possible decomposition routes involve either of the following pairs:



or



These studies are being continued for a more complete elucidation of the decomposition.

### Stability of Florox in Hydrogen Fluoride

At least two instances of nonexplosive decomposition of ClF<sub>3</sub>O have occurred at Rocketdyne in loading stainless-steel lines which, except for possible HF contamination, were considered passive to ClF<sub>3</sub>O. Therefore, the possibility of hydrogen fluoride catalyzed decomposition

CONFIDENTIAL

CONFIDENTIAL



ROCKETDYNE • A DIVISION OF NORTH AMERICAN AVIATION, INC.

or reaction with the metal was examined by adding substantially anhydrous HF to  $\text{ClF}_3\text{O}$  in stainless steel and in Kel-F. No loss of  $\text{ClF}_3\text{O}$  was observed by infrared analysis even with a ten-fold excess of HF at room temperature. Consequently, the observed decompositions have been attributed to nonpassive line connections.

The nonreaction of  $\text{ClF}_3\text{O}$  in HF is noteworthy in that the protonation of the O atom in  $\text{ClF}_3\text{O}$  seemingly does not occur.



This conclusion is based on previous studies in these laboratories in which it was demonstrated that hydroxyl groups were rapidly attacked by  $\text{ClF}_3$ . It is similarly reasoned that a protonated Florox would be highly reactive.

#### The Nature of the Florox-Nitrosyl Fluoride Complex

The possibility of utilizing  $\text{ClF}_3\text{O}$  in forming solid oxidizers with high-energy, basic reagents such as nitrosyl fluoride, suggested investigation of the  $\text{ClF}_3\text{O}$  complex with fluoride bases. The acidic nature of  $\text{ClF}_3\text{O}$  has already been investigated with some alkali metal fluorides and nitrogen oxyfluorides. The mode of complexing and the ionic nature of these complexes are of interest both from a fundamental point of view as well as having a possible utility in the synthesis of solid oxidizers.

The vapor pressure-temperature behavior of a 1:1  $\text{FNO-ClF}_3\text{O}$  mixture indicated a weak complex with a heat of reaction of about -5 kcal from the liquid reagents (Ref. 1). The  $\text{F}^{19}$  n.m.r. of the  $\text{FNO-ClF}_3\text{O}$  system has been

CONFIDENTIAL



CONFIDENTIAL

investigated at low temperatures in hopes of finding evidence for the  $\text{ClF}_4\text{O}^-$  ion. Addition of a fluoride to  $\text{ClF}_3\text{O}$  would be expected to shift the  $\text{F}^{19}$  resonance to higher fields because of increased shielding. Instead, a broad line was observed at 121 ppm relative to fluorine. This position is approximately that predicted because of exchange between  $\text{FNO}$  (-61 ppm) and  $\text{ClF}_3\text{O}$  (150 ppm) in a 1:2 ratio. A weak resonance at 86 ppm was attributed to a  $\text{FClO}_2$  impurity. Some exchange with  $\text{FNO}$  may be responsible for its downfield shift from its previously assigned position at 95 ppm. No resonances attributed to either  $\text{FNO}$  or  $\text{ClF}_3\text{O}$  were found. The results are summarized in Table 3.

TABLE 3

 $\text{F}^{19}$  N.M.R. SPECTRA OF OXYCHLORINE TRIFLUORIDE-NITROSYL FLUORIDE

Complex	Temperature, C	Chemical Shift* ( $\text{F}_2 \approx 0$ ), ppm	Assignment
$\text{FNO}-\text{ClF}_3\text{O}$ ~ 1:1	-77	121 86 (minor)	$\text{FNO}-\text{ClF}_3\text{O}$ exchange $\text{FClO}_2$
$\text{FNO}-\text{ClF}_3\text{O}$ ~ 1:1	26	124 86 (minor)	$\text{FNO}-\text{ClF}_3\text{O}$ exchange $\text{FClO}_2$
$\text{ClF}_3\text{O}$ (neat)	-60 to -70 (slow melting)	150	
$\text{FNO}$	-80	-61	
$\text{FClO}_2$	-60 to -80	95	

\*Shifts were observed relative to an external  $\text{CFCl}_3$  standard and recalculated for  $\delta = 414$  to obtain shifts relative to  $\text{F}_2$ .

CONFIDENTIAL



CONFIDENTIAL



ROCKETDYNE • A DIVISION OF NORTH AMERICAN AVIATION INC

A low-temperature infrared investigation of the  $\text{FNO}-\text{ClF}_3\text{O}$  system has been carried out as well. Although the solid complex may be of the form such as  $\text{ON} \begin{smallmatrix} \text{O} \\ \diagup \text{F} \diagdown \end{smallmatrix} \text{ClF}_3$ , the evidence obtained more clearly supports a molecular complex involving a fluorine bridge.

On becoming solid, the NO stretching frequency in FNO and the ClO stretching frequency in  $\text{ClF}_3\text{O}$  each increased as seen in Table 4. This suggested that contributions such as  $\text{NO}^+\text{F}^-$  and  $\text{ClF}_2\text{O}^+\text{F}^-$  were increased slightly. On formation of the solid  $\text{ClF}_3\text{O}-\text{FNO}$  complex, the ClO frequency was somewhat reduced while the NO frequency was increased and broadened. These observations suggested a partial transfer of the fluoride of nitrosyl fluoride to  $\text{ClF}_3\text{O}$  thus increasing the NO bond order while reducing that of the ClO bond. No meaningful interpretation of ClF or NF infrared data could be made. The N-F solid band was either past the range of the instrument (15.0 microns) or too weak to be observed.

A measure of the relative acidity of  $\text{ClF}_3\text{O}$  was estimated by the extent of FNO fluoride transfer. An acidity range for materials relative to solid FNO was roughly defined by  $\text{NO}^+\text{SbF}_6^-$  at one end,  $\text{NO} = 2385 \text{ cm}^{-1}$ , and FNO solid at the other,  $\text{NO} = 1980 \text{ cm}^{-1}$  (Ref. 3).

On the basis of the hypochromic shift of the NO frequency in the  $\text{FNO}-\text{ClF}_3\text{O}$  complex,  $\text{ClF}_3\text{O}$  was estimated to be about 15 percent as effective an acid as  $\text{SbF}_5$  toward FNO. The low ionic character of the complex  $\text{ClF}_3\text{O}-\text{FNO}$  does not make similar adducts such as  $\text{ClF}_3\text{O}-\text{FNO}_2$  or  $\text{ClF}_3\text{O}-\text{NF}_3\text{O}$  promising as stable solid oxidizer formulations because the expected basic strengths of  $\text{FNO}_2$  and  $\text{NF}_3\text{O}$  are less than that of FNO.

#### Fluorination of Calcium Hypochlorite

As a possibly more convenient route to  $\text{ClF}_3\text{O}$ , the low-temperature fluorination of 95-percent calcium hypochlorite, has been attempted. Also, the

CONFIDENTIAL



ROCKETDYNE • A DIVISION OF NORTH AMERICAN AVIATION INC

CONFIDENTIAL

TABLE 4

SELECTED INFRARED FEATURES IN THE  $\text{FNO-ClF}_3\text{O}$  SYSTEM

Sample	Temperature, C	Vibration, $\text{cm}^{-1}$			
		ClF	ClO	NO	NF
$\text{ClF}_3\text{O}$	Ambient	674	1225	--	--
FNO	Ambient	--	--	1850	765
$\text{ClF}_3\text{O-FNO}$	Ambient	670	1225	1850	765
$\text{ClF}_3\text{O}$	-196	685	1250	--	--
FNO	-196	--	--	1990	?
$\text{FNO-ClF}_3\text{O}$	-196	Broad	1230	2050	?

CONFIDENTIAL

# CONFIDENTIAL




ROCKETDYNE • A DIVISION OF NORTH AMERICAN AVIATION, INC.



search for chlorosyl fluoride was simultaneously pursued. Excess fluorine converted calcium hypochlorite to approximately equal amounts of  $\text{FClO}_2$  and  $\text{ClF}_3$  at  $-80^\circ\text{C}$  with smaller amounts of  $\text{FClO}_3$ . Similar results were obtained with lithium hypochlorite by workers at Picatinny Arsenal (Ref. 4). In the presence of  $\text{KF}$ , only  $\text{FClO}_2$  and  $\text{FClO}_3$  were observed; the  $\text{ClF}_3$  was presumably tied up as the potassium fluoride complex. When limiting amounts of fluorine were used instead, the products were  $\text{FClO}_2$  and  $\text{ClF}_3$ .

It is of interest that the covalent hypochlorites  $\text{ClOCl}$  and  $\text{ClONO}_2$  and  $\text{ClF}_3\text{O}$  upon fluorination while the ionic species,  $\text{Ca}(\text{OCl})_2$ , fails to react to the same products. These contrasting results suggest not only different mechanisms of fluorination but also that the "neutralization" of charge on an anionic substrate is a primary process in a fluorination reaction.

If we consider the  $\text{OCl}^-$  ion in a fluorination reaction, the primary step in its reaction with  $\text{F}^\bullet$  can be one of two reactions; i.e.



When the initial step is as in Eq. 10 or a "neutralization," the resulting short-lived  $\text{OCl}$  may rapidly decompose to  $\text{Cl}_2$  and  $\text{O}_2$ . On the other hand, the initial oxidation of the chlorine atom (Eq. 11) always yields a stable anion. Perhaps the concept of charge neutralization can be further tested by examination of the fluorination of  $\text{KClF}_2$  under flow conditions at temperatures below which neither of the following equilibria may be established:



CONFIDENTIAL

CONFIDENTIAL



ROCKETDYNE • A DIVISION OF NORTH AMERICAN AVIATION INC

Such reactions are:



Thus, if the anion  $\text{ClF}_2^-$  is first "neutralized" (Eq. 12) there is a high probability of "liberating"  $\text{ClF}_3$  from the system (the reaction in Eq. 14 would be slow under flow conditions). On the other hand, the oxidation of the central atom without the neutralization step results in the stable chlorotetrafluoride:



#### FLOROX COMPLEXES WITH ALI METAL FLUORIDES

The complex of  $\text{ClF}_3\text{O}$  and  $\text{KF}$  was prepared by reaction at ambient temperature. Although dissociation pressure was too small to be discernible by direct pressure measurement, continuous slow evolution upon pumping yielded 85 percent of the complexed  $\text{ClF}_3\text{O}$  in 19 hours. This complex was, as expected, weaker than the complexes  $\text{KF}-\text{ClF}_3$  and  $\text{CsF}-\text{ClF}_3\text{O}$ .

One of the difficulties associated with the synthesis of Florox from  $\text{Cl}_2\text{O}$ ,  $\text{CsF}$ , and  $\text{F}_2$  is the problem of separating Florox from the by-product  $\text{ClF}_3$  (Ref. 1). While much of the Florox is obtained as a free gas on warming to room temperature, a considerable portion is retained in the solid phase as the  $\text{CsF}$  complex. Unprogrammed heating has previously shown that this additional product is liberated readily but is accompanied by

CONFIDENTIAL

# CONFIDENTIAL




ROCKETDYNE • A DIVISION OF NORTH AMERICAN AVIATION INC

the evolution of  $\text{ClF}_3$ . During this period it has been determined that complexed Florox and  $\text{ClF}_3$  can be separated effectively by careful thermal dissociation. Thus, at 150 C and with pumping, nearly all the Florox was evolved without any  $\text{ClF}_3$ . At 200 C a small additional amount of Florox was liberated, contaminated with trace amounts of  $\text{ClF}_3$ .




## Reduction of Florox

The reduction of  $\text{ClF}_3\text{O}$  with AgF was considered as in the following equation:



Accordingly, Florox was passed over hot silver fluoride at 135 C and the products were quenched immediately thereafter. Partial reaction gave chlorine and chloryl fluoride as the only products.

In other attempts to synthesize FClO by chemical reduction of  $\text{ClF}_3\text{O}$ ,  $\text{Cl}_2$  and  $\text{Cl}_2\text{O}$  were utilized as potential reducing agents. With  $\text{Cl}_2$ , no reaction was observed at ambient temperature with either  $\text{ClF}_3\text{O}$  or its CsF complex. A series of reactions between  $\text{Cl}_2\text{O}$  and both  $\text{ClF}_3\text{O}$  and its CsF complex were run in varying reactant ratios at ambient temperature and at -18 C. In all cases the products were ClF and  $\text{FClO}_2$  in an overall reaction best described by:




CONFIDENTIAL

CONFIDENTIAL



ROCKETDYNE • A DIVISION OF NORTH AMERICAN AVIATION INC

It is possible that the sequence involves formation of the desired  $\text{FClO}$  followed by its disproportionation:




If the intermediate  $\text{FClO}$  was generated, in no case was it stabilized through complex formation with  $\text{CsF}$ .

#### Fluorination of $\text{Cl}_2\text{O}$

Because the observed unknown thought to be  $\text{FClO}$  arises from the pyrolysis of the fluorination products of  $\text{Cl}_2\text{O}$ , it is either formed by the pyrolysis or released from its complex,  $\text{MClF}_2\text{O}$ , by pyrolysis. Because the desired  $\text{FClO}$  may be strongly complexed, even by  $\text{KF}$ , it was decided to explore the synthesis of  $\text{FClO}$  from fluorination of  $\text{Cl}_2\text{O}$  both in the presence of  $\text{NaF}$  and in the absence of any alkali metal fluoride. Instead of producing  $\text{FClO}$ , however, the low-temperature fluorination of  $\text{Cl}_2\text{O}$  in both cases gave good yields (listed elsewhere in this report) of  $\text{ClF}_3\text{O}$ . Therefore, it was decided to examine the action of the milder fluorinating agents,  $\text{ClF}_3$  and  $\text{ClF}_5$ , on alkali fluoride complexed  $\text{Cl}_2\text{O}$ .

In the case of  $\text{ClF}_5$ , reactions were conducted at  $-50$  and  $0^\circ\text{C}$  with the preformed  $\text{CsF}\cdot\text{Cl}_2\text{O}$  complex and excess  $\text{ClF}_5$ . At the lower temperature, incomplete reaction of the  $\text{Cl}_2\text{O}$  was observed but at  $0^\circ\text{C}$ , all the  $\text{Cl}_2\text{O}$  was consumed. No new products were found. The results of the reactions were in good agreement with the following stoichiometry:




CONFIDENTIAL

# CONFIDENTIAL



ROCKETDYNE • A DIVISION OF NORTH AMERICAN AVIATION, INC.

The  $\text{ClF}_3$  fluorinations utilized NaF to complex the  $\text{Cl}_2\text{O}$  because CsF was expected to complex  $\text{ClF}_3$  faster than reaction with  $\text{Cl}_2\text{O}$  would occur. Again, no products of interest were formed. Correcting for the excess  $\text{ClF}_3$ , the complete reaction of  $\text{Cl}_2\text{O}$  agreed closely with the indicated equation:



While both these reactions could be explained by the formation and subsequent disproportionation of  $\text{FClO}$ ;



no direct evidence for this was obtained.

Earlier (Ref. 2) it was found that in situ fluorination of the products from the reaction of  $\text{HgO}$  and  $\text{Cl}_2$  gave traces of an unknown which has been designated Compound C and may be  $\text{FClO}$ . Also formed were  $\text{ClF}$ ,  $\text{ClF}_3$ , and  $\text{ClF}_5$ . Because of the possible deleterious effect on concomitant HF formation to new F, Cl, O compounds, this reaction was repeated using vacuum dried  $\text{HgO}$ . Two reactions at  $-80^\circ \text{C}$  were run and little, if any, HF was formed. The first reaction gave small quantities of Compound C but most of the initial  $\text{Cl}_2$  was recovered. This latter result is attributed to decomposition of most of the  $\text{Cl}_2\text{O}$  prior to the fluorination step. Because only small amounts of the unknown were obtained and also because experience has shown that it is easily decomposed, no separation was attempted. Instead, an indirect proof of the nature of the unknown will be tried. The unknown is contaminated with  $\text{ClF}$  and small amounts


# CONFIDENTIAL

# CONFIDENTIAL



ROCKETDYNE • A DIVISION OF NORTH AMERICAN AVIATION INC

of  $\text{FClO}_2$  and  $\text{ClF}_5$  which will not fluorinate to yield Florox. But Compound C, if it is  $\text{FClO}$ , would be expected to fluorinate readily to  $\text{ClF}_3\text{O}$ .




A test is now in progress.

## ATTEMPTS AT NEW OXYHALOGEN FLUORIDE OXIDIZERS

### Florox and Oxygen

The reaction between oxygen and  $\text{ClF}_3\text{O}$  has been studied both as a possible route to the unknown oxidizer  $\text{ClF}_3\text{O}_2$  and to determine if oxygen contamination yields  $\text{FClO}_2$ .



When  $\text{ClF}_3\text{O}$  (both liquid and vapor) was exposed to  $\text{O}_2$  at temperatures from ambient to  $-196^\circ\text{C}$ , no reaction occurred as demonstrated both by lack of oxygen uptake and by an unchanged infrared spectrum.

### Reaction of $\text{AsF}_5$ and $\text{Cl}_2\text{O}$

As part of the effort to examine various single bonded Cl-O species as possible precursors to oxychlorine fluoride or Florox, an investigation

CONFIDENTIAL






# CONFIDENTIAL



ROCKETDYNE • A DIVISION OF NORTH AMERICAN AVIATION INC

of the reported compound,  $\text{ClO} \cdot \text{AsF}_5$  (Ref. 5) was initiated. The reactions reported in the literature, together with the proposed fluorinations, are as follows:



At the outset of this work, it was noted that there is a literature discrepancy with regard to the infrared spectrum of  $\text{AsF}_5$ . Samples of  $\text{AsF}_5$  (Ozark-Mahoning) gave an infrared spectrum nearly identical with that reported for " $\text{AsOF}_3$ " rather than  $\text{AsF}_5$  (Ref. 6). But these vendor samples also gave the same infrared spectrum as that obtained for  $\text{AsF}_5$  according to the unpublished thesis of L. K. Akers (Ref. 7). To establish the character of the supplied material, a vapor phase molecular weight determination was made. This gave a value of 169.7 grams/mole vs 169.9 for  $\text{AsF}_5$  and 147.9 for  $\text{AsOF}_3$ . The mass spectrum of the material showed it to be 90-percent  $\text{AsF}_5$  with approximately 10-percent As, O, F species. Because a vapor phase chromatogram showed only one component, it appears the sample was pure  $\text{AsF}_5$ . The As, O, and F impurities undoubtedly arose through reactions of the  $\text{AsF}_5$  with an incompletely dry glass inlet system of the mass spectrometer, because HF and  $\text{SiF}_4$  were also found in the mass spectrum.

Thus, the infrared spectrum reported by Akers (Ref. 7) is correct. Mitra's spectrum for " $\text{AsOF}_3$ " (Ref. 6) consists of  $\text{AsF}_5$  and the background produced on NaCl infrared cell windows after contact with  $\text{AsF}_5$ . Finally, Mitra's

CONFIDENTIAL

CONFIDENTIAL



ROCKETDYNE • A DIVISION OF NORTH AMERICAN AVIATION, INC

infrared spectrum for  $\text{AsF}_5$  is identical in all respects with this window background band only ( $705 \text{ cm}^{-1}$ ). This was shown experimentally and it is probable that this band is attributable to an  $\text{AsF}_6^-$  species. For example,  $\text{K} + \text{AsF}_6^-$  salt (Ref. 8) has its strong band at  $694 \text{ cm}^{-1}$  vs the  $705 \text{ cm}^{-1}$  observed for this peak.

The reaction of  $\text{Cl}_2\text{O}$  and  $\text{AsF}_5$  when examined at  $-80^\circ \text{C}$  did not proceed as indicated in the literature. Mixing the two reactants at  $-196^\circ \text{C}$  and warming to  $-80^\circ \text{C}$  resulted in the formation of a dark red solid which, over a period of several hours, became almost black. Pumping on the sample at this point resulted in the evolution of  $\text{Cl}_2$  (with little or no  $-196^\circ \text{C}$  noncondensables). This  $\text{Cl}_2$  represented most of the Cl in the original  $\text{Cl}_2\text{O}$ . Further warming to room temperature caused additional evolution of small amounts of  $\text{Cl}_2$  and  $\text{ClO}_2$ . When excess  $\text{Cl}_2\text{O}$  was used, no  $\text{AsF}_5$  was recovered in the volatile phase. Remaining at room temperature was a white solid which exhibited two infrared active bands at 7.9 and 14.6 microns. It fumed in moist air and exploded on contact with acid KI. Because the total amount of evolved  $\text{Cl}_2$  was near that contained in the original  $\text{Cl}_2\text{O}$ , the composition of this solid approached  $\text{AsF}_5 \cdot \text{O}$ . Fluorination of the reaction mixture from which only part of the  $\text{Cl}_2$  was removed gave the same white solid product on work-up. Additional characterization work is presently in progress to elucidate the exact character of this new solid material.

#### The Fluorination of $\text{Br}_2\text{O}$

The synthesis of new oxybromine fluorides using  $\text{Br}_2\text{O}$  as a starting material was pursued. This work was based on the analogous, proven

CONFIDENTIAL

# CONFIDENTIAL




ROCKETDYNE • A DIVISION OF NORTH AMERICAN AVIATION, INC.

$\text{Cl}_2\text{O}$  reactions which yielded oxychlorine trifluoride. Fluorination of MF complexed  $\text{Br}_2\text{O}$  at  $-50^\circ\text{C}$  gave as volatile products,  $\text{BrF}_5$ , unreacted  $\text{Br}_2$ , and traces of unstable, unidentified material. While it was anticipated that the sought Br, F, O compounds might remain as complexed solids after the fluorination, pyrolysis up to  $\sim 400^\circ\text{C}$  did not yield any new products. The failure of this reaction may be caused by the inherent instability of the starting material or thermal degradation of the desired products upon pyrolysis.

Because chlorine nitrate has also been shown to be a useful precursor to  $\text{ClF}_3\text{O}$ , it was decided to utilize bromine nitrate,  $\text{BrONO}_2$ , as an intermediate in the preparation of new oxybromine fluorides. This compound offered two advantages over  $\text{Br}_2\text{O}$ . It is more stable than  $\text{Br}_2\text{O}$ , decomposing around  $0^\circ\text{C}$  vs approximately  $-40^\circ\text{C}$  for  $\text{Br}_2\text{O}$ . Also, based on the high yields of Florox obtained by fluorination of uncomplexed  $\text{ClONO}_2$ , the analogous application of  $\text{BrONO}_2$  can be expected to give similar results while eliminating the necessity of complexing the desired products.

The synthesis of  $\text{BrONO}_2$  was conducted according to the reported literature procedure (Ref. 9).



Fluorination of the product from which  $\text{Cl}_2$  and excess  $\text{ClNO}_3$  were removed is in progress at  $-50^\circ\text{C}$ .

# CONFIDENTIAL

CONFIDENTIAL



ROCKETDYNE • A DIVISION OF NORTH AMERICAN AVIATION INC

#### CESIUM FLUORIDE-CHLORINE MONOXIDE COMPLEX STUDIES

Further investigation of the stoichiometry of  $\text{CsF-Cl}_2\text{O}$  complexes has given reproducible ratios in contrast to previously reported experiments (Ref. 1). To determine the maximum ratio  $\text{Cl}_2\text{O/CsF}$  in the complex, formed at  $-80^\circ\text{C}$ , large excesses of  $\text{Cl}_2\text{O}$  were stored over  $\text{CsF}$  for several days and then pumped on overnight at  $-80^\circ\text{C}$  to remove the uncomplexed  $\text{Cl}_2\text{O}$ . Experimental  $\text{Cl}_2\text{O/CsF}$  ratios of 1.5, 1.42, 1.48, and 1.54 were obtained. In another run, pumping at  $-80^\circ\text{C}$  for 3-1/2 days resulted in a 1.49  $\text{Cl}_2\text{O/CsF}$  stoichiometry. Only 0.17  $\text{Cl}_2\text{O}$  per  $\text{CsF}$  was removed after overnight pumping. It is apparent that a slow forming complex with a  $\text{Cl}_2\text{O/CsF}$  ratio at or near 1.5 is formed at  $-80^\circ\text{C}$ . A sample of the complex was exposed to the air, hit with a hammer, and heated with a torch with no explosive results.

CONFIDENTIAL

CONFIDENTIAL



ROCKETDYNE • A DIVISION OF NORTH AMERICAN AVIATION INC

#### EXPERIMENTAL EFFORT

##### LOW-TEMPERATURE INFRARED CELL

The low-temperature infrared experiments were determined in a simple, infrared cell which was constructed as described elsewhere (Ref. 10). The completed assembled cell fits conveniently into a Perkin-Elmer 137 Infracord spectrometer and has a cooling capacity of 2 liters. It can be used at temperatures as low as 77 K and is relatively easy to use with slush baths because the cooling flask is insulated with 3-inch solid foam. The inner window of AgCl fits into a copper optical blank holder. The body is glass and is fitted with two outside AgCl windows.

Florox complexes were preformed in a metal vacuum line and condensed onto the inner AgCl window at -196 C by means of a copper entrance tube directed at the window.

##### SYNTHESIS OF FLOROX

Oxychlorine trifluoride,  $\text{ClF}_3\text{O}$ , was synthesized by fluorination of  $\text{Cl}_2\text{O}$  at -80 C in the presence and absence of added alkali metal fluoride. In most cases the reaction was conducted in a 300-milliliter stainless-steel cylinder previously passivated by exposure to at least one atmosphere of fluorine for a period of at least 16 hours. Chlorine monoxide and fluorine were introduced into the reactor by distillation in vacuo. Separation of the  $\text{ClF}_3\text{O}$  product from side products was achieved in all cases by fractional condensation.

CONFIDENTIAL

CONFIDENTIAL



ROCKETDYNE • A DIVISION OF NORTH AMERICAN AVIATION INC

Excess fluorine and any produced oxygen passed  $-196^{\circ}\text{C}$ . The next most volatile side products were  $\text{FClO}_2$ ,  $\text{ClF}$ , and  $\text{Cl}_2$  and were removed by passage through a cold trap at  $-95^{\circ}\text{C}$  with the  $\text{ClF}_3\text{O}$  being retained. Chlorine trifluoride was partially retained at  $-95^{\circ}\text{C}$ ; therefore, removal of  $\text{ClF}_3$  was achieved by repeated passage through a trap held at  $-80^{\circ}\text{C}$ , with some loss of  $\text{ClF}_3\text{O}$ .

CONFIDENTIAL

CONFIDENTIAL



ROCKETDYNE • A DIVISION OF NORTH AMERICAN AVIATION, INC

REFERENCES

1. R-5883-5, Inorganic Halogen Oxidizers, Rocketdyne, a Division of North American Aviation, Inc., Canoga Park, California, 30 December 1965, CONFIDENTIAL.
2. R-5883-4, Inorganic Halogen Oxidizers, Rocketdyne, a Division of North American Aviation, Inc., Canoga Park, California, 30 September 1965, CONFIDENTIAL.
3. Sharp, D. W. A. and J. Thorley, J. Chem. Soc., 3557 (1963).
4. TR 3285, Oxidizer Chemistry, AMCMS 5221.11.145.02, Liquid Rocket Propulsion Laboratory, U. S. Army Munitions Command, Picatinny Arsenal, Dover, New Jersey, November 1965, CONFIDENTIAL.
5. Schmeisser, M. and W. Fink, Angew. Chem., 69, 780 (1957).
6. Mitra, G., J. Am. Chem. Soc., 80, 5639 (1958).
7. Akers, L. K., Ph.D. Thesis, Vanderbilt University, 1955.
8. Pitts, J. J. et al, Inorg. Chem., 4, 257 (1965).
9. Schmeisser, M. and L. Taglinger, Chem. Ber., 94, 1533 (1961).
10. R-5785, Annual Summary Report, Research in High-Energy Oxidizers, Rocketdyne, a Division of North American Aviation, Inc., Canoga Park, California, August 1964, CONFIDENTIAL.

CONFIDENTIAL

CONFIDENTIAL

Security Classification

DOCUMENT CONTROL DATA - R&D		
(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)		
1 ORIGINATING ACTIVITY (Corporate author)		2a REPORT SECURITY CLASSIFICATION
Rocketdyne, a Division of North American Aviation, Inc., 6633 Canoga Ave., Canoga Park, California		CONFIDENTIAL
		2b GROUP
		1
3 REPORT TITLE		
INORGANIC HALOGEN OXIDIZERS		
4 DESCRIPTIVE NOTES (Type of report and inclusive dates)		
Quarterly (29 November 1965 through 28 February 1966)		
5 AUTHOR(S) (Last name, first name, initial)		
Pilipovich, D.; Bauer, H. F.; Lindahl, C. B.; Schack, C. J.		
6 REPORT DATE	7a TOTAL NO OF PAGES	7b NO OF REFS
30 March 1966	27	10
8a CONTRACT OR GRANT NO.	9a ORIGINATOR'S REPORT NUMBER(S)	
Nonr 4428(00)	R-5883-6	
b PROJECT NO.		
c ARPA Order No. 23	9b OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
d		
10 AVAILABILITY/LIMITATION NOTICES		
11 SUPPLEMENTARY NOTES	12 SPONSORING MILITARY ACTIVITY	
	Advanced Research Projects Agency Washington 25, D.C.	
13 ABSTRACT Oxychlorine trifluoride has been synthesized from $\text{Cl}_2\text{O} + \text{F}_2$ in the presence of $\text{CsF}$ , $\text{RbF}$ , $\text{KF}$ , and $\text{NaF}$ as well as in their absence. Oxychlorine trifluoride has also been synthesized from uncomplexed chlorine nitrate in 84-percent yield. Fluorination in situ of the $\text{Cl}_2 + \text{HgO}$ reaction products has also produced $\text{ClF}_3\text{O}$ . Florox was stable in excess hydrogen fluoride or excess oxygen at ambient temperature and stable to 5-day storage at 71 C in Monel. In a flow system, thermal decomposition of Florox started near 300 C and was complete below 586 C. The predominant product recovered above 300 C was chlorine monofluoride. The complex between Florox and $\text{CsF}$ dissociated below 150 C and allowed complete removal of contaminant $\text{ClF}_3$ as the remaining nonvolatile solid $\text{CsClF}_4$ . Investigation of the FNO complex of $\text{ClF}_3\text{O}$ by $\text{F}^{19}$ n.m.r. showed exchange at -77 C; an infrared study at -196 C indicated the complex to be predominantly covalent. Attempts to reduce $\text{ClF}_3\text{O}$ to $\text{FCIO}$ have been unsuccessfully attempted with $\text{AgF}$ , $\text{Cl}_2$ , and $\text{Cl}_2\text{O}$ . Fluorination of $\text{Cl}_2\text{O}$ by $\text{F}_2$ , $\text{ClF}_3$ , and $\text{ClF}_5$ has also failed to produce $\text{FCIO}$ . As a new route to oxychlorine fluorides, the reported intermediate $\text{ClO} \cdot \text{AsF}_5$ has been investigated. The synthesis of new bromine oxyfluorides from $\text{Br}_2\text{O}$ complexes failed, giving mainly $\text{BrF}_5$ . Bromine nitrate has been prepared as a possible alternate precursor to $\text{BrF}_3\text{O}$ or $\text{BrF}_5\text{O}$ . Although the nature of the bonding in the $\text{Cl}_2\text{O} \cdot \text{CsF}$ complex has not been clarified, a reproducible composition, $\text{CsF} \cdot 1.5 \text{Cl}_2\text{O}$ has been obtained near -80 C. (C)		

DD FORM 1473

JAN 64

CONFIDENTIAL  
Security Classification



CONFIDENTIAL

Security Classification

14 KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
Chlorines						
Fluorines						
Fluorination						
Stoichiometry						
Hydrolysis						
Glow Discharges						

# INSTRUCTIONS

1. **ORIGINATING ACTIVITY:** Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (*corporate author*) issuing the report.

2a. **REPORT SECURITY CLASSIFICATION:** Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.

2b. **GROUP:** Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.

3. **REPORT TITLE:** Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.

4. **DESCRIPTIVE NOTES:** If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.

5. **AUTHOR(S):** Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.

6. **REPORT DATE:** Enter the date of the report as day, month, year; or month, year. If more than one date appears on the report, use date of publication.

7a. **TOTAL NUMBER OF PAGES:** The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.

7b. **NUMBER OF REFERENCES:** Enter the total number of references cited in the report.

8a. **CONTRACT OR GRANT NUMBER:** If appropriate, enter the applicable number of the contract or grant under which the report was written.

8b, 8c, & 8d. **PROJECT NUMBER:** Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.

9a. **ORIGINATOR'S REPORT NUMBER(S):** Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.

9b. **OTHER REPORT NUMBER(S):** If the report has been assigned any other report numbers (*either by the originator or by the sponsor*), also enter this number(s).

10. **AVAILABILITY/LIMITATION NOTICES:** Enter any limitations on further dissemination of the report, other than those

imposed by security classification, using standard statements such as:

- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through \_\_\_\_\_."
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through \_\_\_\_\_."
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through \_\_\_\_\_."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

11. **SUPPLEMENTARY NOTES:** Use for additional explanatory notes.

12. **SPONSORING MILITARY ACTIVITY:** Enter the name of the departmental project office or laboratory sponsoring (*paying for*) the research and development. Include address.

13. **ABSTRACT:** Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. **KEY WORDS:** Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical content. The assignment of links, rules, and weights is optional.

CONFIDENTIAL

Security Classification